


Serial No. 09/937,912  
Art Unit: 1751

Docket No. H3933 PCT/US


In the Claims

1-13. (Canceled)

14. (Currently amended) A composition for coloring keratin fibers comprising  
(a) at least one tenside of formula (I)



wherein y is an integer from 0 to 2, x is an integer from 1 to 3, and the sum of x and y is 3, wherein M is hydrogen, an alkali metal, alkaline earth metal, or an ammonium cation, or an alkyl radical having 1 to 4 carbon atoms that is optionally substituted by one or more hydroxyl groups, wherein B is a physiologically compatible anion, and wherein R is a radical of formula (II),



in which z is an integer from 1 to 4, R<sup>1</sup> and R<sup>2</sup>, independently of one another, are a C<sub>1</sub> to C<sub>4</sub> alkyl radical, that is optionally substituted by one or more hydroxyl groups, or an acyl group, A is -O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -O-CH<sub>2</sub>-CH<sub>2</sub>- or -O-CH<sub>2</sub>-CHOH-CH<sub>2</sub>-, and R<sup>3</sup> is a branched or unbranched, saturated C<sub>8</sub> to C<sub>18</sub> acyl radical, or a branched or unbranched, monounsaturated or polyunsaturated C<sub>8</sub> to C<sub>18</sub> acyl radical;

(b) at least one conditioning component; and

Serial No. 09/937,912  
Art Unit: 1751

Docket No. H3933 PCT/US

- (c) at least one dye or dye precursor, or combinations thereof; and  
(d) at least one anionic tenside.

15. (Canceled)
16. (Currently amended) The composition of claim 14 ~~15~~, wherein the anionic tenside comprises a soap.
17. (Previously presented) The composition of claim 14 wherein the conditioning component comprises a low molecular weight quaternary ammonium compound.
18. (Previously presented) The composition of claim 14 wherein the conditioning component comprises a cationic polymer.
19. (Previously presented) The composition of claim 18 wherein the cationic polymer comprises a quaternized cellulose derivative.
20. (Previously presented) The composition of claim 18 wherein the cationic polymer comprises Polyquaternium-2.
21. (Previously presented) The composition of claim 14 wherein the conditioning component comprises a quaternized protein hydrolyzate.
22. (Previously presented) The composition of claim 14 wherein the conditioning component comprises a silicone oil.
23. (Previously presented) The composition of claim 14 wherein the dye or dye precursor

Serial No. 09/937,912  
Art Unit: 1751

Docket No. H3933 PCT/US

comprises at least one oxidative developer dye precursor.

24. (Previously presented) The composition of claim 14 wherein the dye or dye precursor comprises at least one indole derivative, or indoline derivative, or combinations thereof.


25. (Previously presented) The composition of claim 14 wherein the dye or dye precursor comprises at least one substantive dye, or natural dye, or combinations thereof.

26. (Previously presented) The composition of claim 14 wherein the tenside of formula I comprises at least one compound selected from Linoleamidopropyl PG-Dimonium Chloride Phosphate, Cocamidopropyl PG-Dimonium Chloride Phosphate or Stearamidopropyl PG-Dimonium Chloride Phosphate, or combinations thereof.

27. (Previously presented) The composition of claim 26 wherein the conditioning component comprises at least one low molecular weight quaternary ammonium compound or cationic polymer, or combinations thereof.

28. (Previously presented) A method for coloring keratin fibers comprising applying to keratin fibers a composition comprising


(a) at least one tenside of formula (I)



Serial No. 09/937,912  
Art Unit: 1751

Docket No. H3933 PCT/US

wherein y is an integer from 0 to 2, x is an integer from 1 to 3, and the sum of x and y is 3, wherein M is hydrogen, an alkali metal, alkaline earth metal, or an ammonium cation, or an alkyl radical having 1 to 4 carbon atoms that is optionally substituted by one or more hydroxyl groups, wherein B is a physiologically compatible anion, and wherein R is a radical of formula (II),



in which z is an integer from 1 to 4, R<sup>1</sup> and R<sup>2</sup>, independently of one another, are a C<sub>1</sub> to C<sub>4</sub> alkyl radical, that is optionally substituted by one or more hydroxyl groups, or an acyl group, A is -O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -O-CH<sub>2</sub>-CH<sub>2</sub>- or -O-CH<sub>2</sub>-CHOH-CH<sub>2</sub>-, and R<sup>3</sup> is a branched or unbranched, saturated C<sub>8</sub> to C<sub>18</sub> acyl radical, or a branched or unbranched, monounsaturated or polyunsaturated C<sub>8</sub> to C<sub>18</sub> acyl radical;

(b) at least one conditioning component; and

(c) at least one dye or dye precursor, or combinations thereof.

29. (Previously presented) The method of claim 28 wherein the composition further comprises at least one anionic tenside.

30. (Previously presented) The method of claim 29 wherein the conditioning component comprises at least one low molecular weight quaternary ammonium compound, or cationic polymer, or combinations thereof.

31. (Previously presented) The method of claim 30 wherein the tenside of formula I comprises at least one compound selected from Linoleamidopropyl PG-Dimonium Chloride

**Serial No. 09/937,912**  
**Art Unit: 1751**

**Docket No. H3933 PCT/US**

Phosphate, Cocamidopropyl PG-Dimonium Chloride Phosphate or Stearamidopropyl PG-Dimonium Chloride Phosphate, or combinations thereof.

32. (Previously presented) The method of claim 30 wherein the anionic tenside comprises a soap.